

## Harnessing the Power of Neuroplasticity to Restore Physical Function

June 7, 2020

By: Garrett Salpeter, Founder of NeuFit

## **Scientific References**

- 1. Lang, C.E., et. al. *Observation of amounts of movement practice provided during stroke rehabilitation*. Arch Phys Med Rehabil 90(10): 1692-1698, 2009
- 2. Tommassini, V., et. al., *Neuroplasticity and Functional Recovery in Multiple Sclerosis*. Nat Rev Neurol. 8(11): 635-646, 2012
- 3. Irvine, K.A., Blakemore W.F., *Remyelination protects axons from demyelination-associated axon degeneration.* Brain 131: 1464-1477, 2008
- 4. Hebb, D.O. *Organization of behavior*. Wiley; New York: 1949
- 5. Cramer S.C., Repairing the human brain after stroke: Mechanisms of spontaneous recovery. Ann. Neurol. 62:272-287, 2008
- 6. Fridman, E.A., et. al., *Reorganization of the human ipsilesional premotor cortex after stroke.* Brain. 127: 747-758, 2004
- 7. Jacobs, K.M., et. al. Reshaping the cortical motor map by unmasking latent intracortical connections. Science 251: 944-947, 1991
- 8. Pineiro, R., et. al. *Functional MRI detects posterior shfits in primary sensorimotor cortex activation after stroke: evidence of local adaptive reorganization?* Stroke 32: 1134-1139, 2001
- 9. Nudo, R.J., *Use-dependent alterations of movement representations in primary motor cortex of adult squirrel monkeys.* J. Neurosci 16(2): 785-807, 1996
- 10. Adkins, D.L., et. al. *Motor training induces experience-specific patterns of plasticity across motor cortex and spinal cord.* J Appl Physiol 101(6): 1776-1782, 2006
- 11. Rocca, M.A., et. al. *Pyramidal tract lesions and movement-associated cortical recruitment in patients with MS.* Neuroimage 23:141-147, 2004
- 12. Johansen-Berg, H., et. al. *Correlation between motor improvements and altered fMRI activity after rehabilitative therapy.* Brain. 125: 2731-2742, 2002
- 13. Lee, M., et. al. *The motor cortex shows adaptive functional changes to brain injury from multiple sclerosis.* Ann Neurol. 47: 606-613, 2000



- 14. Tomassini, V. *Relating Brain Damage to Brain Plasticity in Patients with Multiple Sclerosis*. Neurorehabil Neural Repair. 2012.
- 15. Zeller, D., et. al. *Rapid-onset central motor plasticity in multiple sclerosis.* Neurology 74(9): 728-735, 2010
- 16. Toosy, A.T., et. al. *Adaptive cortical plasticity in higher visual areas after acute optic neuritis*. Ann. Neurol. 57: 622-633, 2005
- 17. Werring, D.J., et. al. Recovery from optic neuritis is associated with a change in the distribution of cerebral response to visual stimulation: a functional magnetic resonance imaging study. J. Neurol Neurosurg Psychiatry. 68: 441-449, 2000
- 18. Pantano, P., et. al. *Cortical motor reorganization after a single clinical attack of multiple sclerosis.* Brain: a journal of neurology. 125: 1607-1615, 2002
- 19. Rocca, M.A., et. al. *Cortical adaptation in patients with MS: a cross-sectional functional MRI study of disease phenotypes.* Lancet Neurol 4: 618-626, 2005
- 20. Rocca, M.A., et. al. Adaptive functional changes in the cerebral cortex of patients with nondisabling multiple sclerosis correlate with the extent of brain structural damage. Ann Neurol 51: 330-339, 2002
- 21. Chiaravalloti, N., et. al. *Cerebral activation patterns during working memory performance in multiple sclerosis using FMRI.* J Clin Exp Neuropsychol. 27: 33-54, 2005
- 22. Mainero, C., et. al. *FMRI evidence of brain reorganization during attention and memory tasks in multiple sclerosis.* Neuroimage. 21: 858-867, 2004
- 23. Reddy, H., et. al. *Functional brain reorganization for hand movement in patients with multiple sclerosis: defining distinct effects of injury and disability.* Brain. 125: 2646-2657, 2002
- 24. Taub, E., et. al. An operant approach to rehabilitation medicine: overcoming learned nonuse by shaping. *J Exp Anal Behav. 61(2):281-93, 1994*
- 25. Sumowski, J.F., Wylie, G.R., Deluca, J., Chiaravalloti, N., *Intellectual enrichment is linked to cerebral efficiency in multiple sclerosis: functional magnetic resonance imaging evidene for cognitive reserve.* Brain: a journal of neurology. 133: 362-374, 2010
- 26. Tomassini V., et. al. Sex hormones modulate brain damage in multiple sclerosis: MRI evidence. J Neurol Neurosurg Psychiatry. 76: 272-275, 2005



- 27. Bove, R., Chitnis, T. *The role of gender and sex hormones in determining the onset and outcome of multiple sclerosis.* Mult Scler 20(5): 520-526, 2014
- 28. Gregg, C., et. al. White matter plasticity and enhanced remyelination in the maternal CNS. J Neurosci 27: 1812-1823, 2007
- 29. Reddy, H., et. al. *Relating axonal injury to functional recovery in MS.* Neurology 54(1): 236-239, 2000
- 30. Sterr, A., et. al. *The role of corticospinal tract damage in chronic motor recovery and neurorehabilitation: a pilot study.* Neurorehabil Neural Repair. 24(5):413-9, 2010.
- 31. Nudo, R.J. *Adaptive plasticity in motor cortex: implications for rehabilitation after brain injury.* J Rehabil Med. 2003: 7-10
- 32. Veerbeek, J.M., et. al. What is the evidence for physical therapy poststroke? A systematic review and meta-analysis. PLoS One 9(2):e87987, 2014
- 33. Swain, R.A., et. al. *Prolonged exercise induces angiogenesis and increases cerebral blood volume in primary motor cortex of the rat.* Neuroscience 117(4): 1037-1046, 2003
- 34. Pohl, M., et. al. Repetitive locomotor training and physiotherapy improve walking and basic activities of daily living after stroke: a single-blind, randomized multicentre trial (DEutsche GAngtrainerStudie, DEGAS). Clin Rehabil 21:17–27, 2007
- 35. Tomassini, V., et. al. *Relating brain damage to brain plasticity in patients with multiple sclerosis.* Neurorehabil Neural Repair. 2012
- 36. Mark, V.W., et. al. Constraint-Induced Movement therapy can improve hemiparetic progressive multiple sclerosis. Preliminary findings. Mult Scler. 14(7):992-4, 2008
- 37. Taub, E., et. al. *Technique to improve chronic motor deficit after stroke*. Arch Phys Med Rehabil. 74(4):347-54, 1993
- 38. Uswatte, G., Taub E. *Constraint-induced movement therapy: a method for harnessing neuroplasticity to treat motor disorders.* Prog Brain Res 207: 379-401, 2013
- 39. Mark, V.W., et. al. *Constraint-induced movement therapy for the lower extremities in multiple sclerosis: case series with 4-year follow-up.* Arch Phys Med Rehabil 94(4):753-760, 2013



- 40. Weiss, P.L, et. al. *Virtual reality in neurorehabilitation*. Textbook of Neural Repair and Rehabilitation 51:182-198, 2006
- 41. Pascual-Leone, A., Study and modulation of human cortical excitability with transcranial magnetic stimulation. J Clin Neurophysiol 15(4):333-343, 1998
- 42. Koch, G., et. al. *Improvement of hand dexterity following motor cortex rTMS in multiple sclerosis patients with cerebellar impairment.* Mult Scler 14(7):995-998, 2008
- 43. Nitsche, M.A., Paulus, W. *Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation.* J Physiol 527(3):633-639, 2000
- 44. Cuypers, K., et. al. *Anodal tDCS increases corticospinal output and projection strength in multiple sclerosis.* Neurosci Lett 554:151-155, 2013
- 45. Fritsch, B., et. al. *Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning.* Neuron 66(2):198-204, 2010
- 46. Sampson, P., Using functional electrical stimulation mediated by iterative learning control and robotics to improve arm movement for people with multiple sclerosis. IEEE Trans Neural Syst Rehabil Eng 24(2):235-248, 2016
- 47. Beer, S., et. al. *Robot-assisted gait training in multiple sclerosis: a pilot randomized trial.* Mult Scler 14(2):231-236, 2008
- 48. Min S.S., et. al. *Chronic brain inflammation impairs two forms of long-term potentiation in the rat hippocampal CA1 area.* Neurosci Lett. 456: 20-24, 2009